

## Isolation, characterization and primary screening of soil actinomycetes from Kenyatta University arboretum grounds for antibacterial activities

Osborn Yamakhakha Khasabuli and Anthony Kebira Nyamache

Kenyatta University (main campus), P.O. Box 43844, Nairobi, Telephone: Tel: +254 20 8710901, +254 20 8711622, Fax: +254 20 8711575, Website: [www.ku.ac.ke](http://www.ku.ac.ke) . Email: [info@ku.ac.ke](mailto:info@ku.ac.ke), [Khasabuli10@gmail.com](mailto:Khasabuli10@gmail.com) and [akibera2000@gmail.com](mailto:akibera2000@gmail.com)

Original submitted in on 9<sup>th</sup> November 2012 Published online at [www.m.elewa.org](http://www.m.elewa.org) on 28<sup>th</sup> February 2014.  
<http://dx.doi.org/10.4314/jab.v74i1.12>

### ABSTRACT

*Objective* : Antimicrobial resistance has increased drastically in recent years in the developing countries, and it has leading public health concern. With almost all organisms becoming multidrug resistant to the commonly used antibiotics, there is a need to search for more drugs that are novel in order to address this challenge. Therefore, this study was aimed at isolating antimicrobial producing actinomycetes and to evaluate their secondary metabolites against selected multidrug resistant bacteria isolates.

*Methodology*: Half kg soil samples were collected 15cm below the ground. Starch Casein Agar (SCA) was used to isolate actinomycetes. Fifteen different actinomycetes identified as Is.1-Is.15 were isolated. They were differentiated based on the difference in appearance of the colony morphology and mycelial structure. Their metabolites were tested for antibiotic activities through the primary screening using the disk diffusion methods. Test bacteria were; *Bacillus subtilis*, *E.coli*, *Klebsiella Pneumoniae*, *Staphylococcus aureus*, *Shigella sonei* and *Salmonella typhi*. All the secondary metabolites from the isolated actinomycetes proved to possess the ability to inhibit bacterial growth by formation of different sized halo zones.

*Conclusion and application*: Secondary metabolites produced by actinomycetes have the ability to inhibit growth of pathogenic bacteria. The ability of these metabolites to inhibit growth of test pathogenic bacteria has been attributed to the establishment of the fact that novel antibiotics can be manufactured from metabolites produced by actinomycetes. Large scale manufacture can be done through purification of these metabolites to ensure large scale production of novel drugs. This can serve help clear pathogenic bacteria that are ever developing resistance on the available antibiotics.

Key words: Actinomycetes, metabolites, soil, antimicrobial activity

### INTRODUCTION

Actinomycetes are a group of organisms originally considered to be intermediates between fungi and bacteria. The study of Cummins and Harris (1956) established that actinomycetes have a cell wall composition resembling that of Gram positive bacteria. They are also branching unicellular

microorganisms. They produce mycelium which may either be aerial or may spread on the substrate on which the microorganism is growing (Harvey, 2007). The mycelia in some species may break to form rod- or coccoid-shaped forms. Many genera also form spores; the sporangia or spore cases may be found

on aerial hyphae, on the colony surface or free within the environment. They possess DNA that contains high amounts of Guanine + Cytosine (>55%) (Cummins and Harris, 1956). Among these groups of microorganisms, *Streptomyces* are dominant. The others referred to as the non-actinomycetes are called the rare actinomycetes. The majority of the actinomycetes are free living (Ceylan, Okmen and Ugur, 2008). Actinomycetes are one of the major constituents of soil microorganisms, together with some bacteria as well as some fungi. Some inhabit marine environments while others groups live in mangrove environments (Kumar, 2001). They play an important role in decomposition of organic materials such as cellulose and chitin and thereby playing a vital part in organic matter turnover and carbon cycle (Williams *et al.*, 1971). Actinomycetes are known to produce diverse metabolites from their complex biochemical processes. Over 23,000 microbial secondary metabolites are known including the beta-lactam antibiotics like penicillin and cephalosporins and some form the shikimic acid like chloramphenicol for example (Barrios-Gonzalez *et al.*, 2003). Approximately, 42% of these metabolites are known to be produced by actinomycetes, 16% by fungi and other bacteria. Among various genera, *Streptomyces*, *Saccharopolyspora*, *Amycolatopsis*,

*Micromonospora* and *Actinoplanes* are the major producers of commercially important biomolecules. These metabolites are released into the environment. They have broad spectrum biological activities such as antibacterial, antifungal, antiviral, antiparasitic, immunosuppressive, antitumor, insecticidal, anti-inflammatory, antioxidant, enzyme inhibitory and diabetogenic (Sacramenti *et al.*, 2004). Those that produce antibacterial and antifungal metabolites may pose several effects on the immediate surrounding and the major one is that of inhibiting other microorganisms from growing or multiplying. These groups of actinomycetes are referred to as the biological antagonistic types. They are of special interest since they are the ones that are exploited and their metabolites used in the manufacture of antibiotics. Microbial metabolites are a source of life saving environments for bacterial and fungal infections. This is because they are used in the manufacture of some of the effective antibiotics including: penicillin, erythromycin, streptomycin, tetracyclines, vancomycin and amphotericin. These microbial natural products are notable not only for their potent therapeutic activities but also for the fact that they frequently pose desirable pharmacokinetic properties required for clinical development (Kekuda *et al.*, 2010).

## METHODOLOGY

**Study site and sample collection:** The soil samples were collected from virgin arboretum grounds of Kenyatta University. This enabled evaluation of a group of antibiotic producing actinomycetes that were inhabitants of that region. Half kilograms of soil samples were collected about 10cm below the ground surface and packed in polythene bags and carried to the Kenyatta University convectional laboratories (BZ6PMS) for analysis. The collected samples had labels identifying the field, sample and date of collection. The fields were labeled 1, 2 and 3. Sample from field 1 had labels 1a and 1b, field 2 had labels 2a and 2b finally field 3 had labels 3a and 3b. The digits 1, 2 and 3 were the codes representing the areas from where the samples were collected. The samples were air dried over the benches for one week. This was done to help reduce the population of Gram negative bacteria (Jeffrey *et al.*, 2008).

**Microbe isolation:** The soil suspension technique described by Oskay *et al.*, (2004) was used. One gram of


the soil sample was measured and mixed with 9ml of sterile distilled water (sdH<sub>2</sub>O). The soil suspension was shaken vigorously under room temperature (25 ± 2°C) on an orbital shaker at 200 rpm for 10 min. 100µl of the soil suspension was pipetted and lawn onto Starch Casein Agar (SCA) (Soluble starch, 10.0 g; Casein hydrolysate, 0.3 g; KNO<sub>3</sub>, 2.0 g NaCl, 2.0 g; K<sub>2</sub>HPO<sub>4</sub>, 2.0 g; MgSO<sub>4</sub>.7H<sub>2</sub>O, 0.05 g; FeSO<sub>4</sub>.7H<sub>2</sub>O, 0.01 g; Agar, 18.0 g; distilled water, 1000 ml (Jeffrey, 2008). The plates were done in duplicates. One set was incubated at room temperature while the other set incubated at 37°C. At 37°C colonies were fully grown after 24 hours while at 25°C colonies grew fully after 7-10 days. Incubation at 37°C was efficient since the temperature was favorable for faster growth of actinomycete as well as serve optimum temperature for bacterial growth.

**Identification and characterization:** A total of 15 isolates were obtained and identified as Is.1, Is.2, Is.3 ... and Is.15 based on their differences in colony morphology

(Table 1). Preliminary identification of the colonies was done based on morphological observations both microscopic and macroscopic. Microscopically, filamentous colonies were sub cultured on fresh nutrient agar and incubated at 37°C to obtain pure cultures.

**Microscopic observations:** A simple stain technique using methylene blue was used. A mass of colony was scooped using a wire loop, spread on a clean slide, air

dried and fixed by passing over a flame twice. The slide was then flooded with methylene blue stain and after 3 seconds the excess stain washed off in running tap water. The slide was then blot dried and observed using a light microscope with ×40 magnification objective (Photograph1, pg13). This was a preliminary identification technique for the actinomycetes.



**Photograph1:** Mycelia of isolate 1 (×400mag)


**Macroscopic observations:** This included observing the morphological characteristics of the colonies. Table1 gives the details of the observations.

**Pure cultures:** The cultures were ascertained as pure only when the culture plate had only one group of the organism growing (Michael *et al.*, 2009). Broth cultures

were then prepared by inoculating the pure cultures in nutrient agar broth and incubating at 37°C for one week. These cultures were sealed with parafilm and kept undisturbed in the fridge for subsequent analysis (Plate 2). Also the culture plates were at all times sealed using the parafilm to avoid mites from invading the plates.

**Table1:** Colony characteristics.

Isolate	Colony characterization.
Is.1	Irregular, moist, undulate margin, glistening surface, translucent, crateriform elevation. Colony size ranged from 2mm-4mm.
Is.2	Irregular in shape with an entire margin, crateriform elevation, opaque, moist and a glistening surface.
Is.3	Irregular in shape, entire margin, flat elevation, moist with a glistening surface, creamish opaque.
Is.4	Irregular, translucent, moist, glistening, lobate with a flat elevation.
Is.5	Moist, opaque, glistening surface, filiform margin, flat elevation with an irregular shape.
Is.6	Irregular, flat, lobate, opaque, glistening, moist.
Is.7	Irregular in shape, moist, translucent, glistening surface, crateriform elevation, lobate margin.
Is.8	Circular in shape, entire margin, crateriform, opaque, moist with a glistening surface.
Is.9	Circular in shape, glistening surface, crateriform elevation, entire margin and moist.
Is.10	Opaque, irregular in shape, moist, lobate margin with crateriform elevation.
Is.11	Irregular in shape with a lobate margin, opaque, raised elevation, whitish in color and appears brittle.
Is.12	Dull colonies, whitish, irregular in shape, moist and umbonate elevation.
Is.13	Translucent irregular colonies, crateriform elevation and moist with a lobate margin.
Is.14	Irregular glistening colonies, crateriform, opaque with a lobate margin and whitish in color.
Is.15	Irregular in shape, opaque and whitish, convex elevation, lobate margin, moist and dull.



**Plate2:** Pure isolate1 colonies

**Preservation of pure cultures:** A sub culture was made but this time the media was Glycerol-arginine agar ( Glycerol 12.5g, Arginine 1.0g, NaCl 1.0g, K<sub>2</sub>HPO<sub>4</sub>1.0g, MgSO<sub>4</sub>.7H<sub>2</sub>O 0.5g, Fe<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub>.6H<sub>2</sub>O 0.01g, CuSO<sub>4</sub>.5H<sub>2</sub>O 0.001g, ZnSO<sub>4</sub>.7H<sub>2</sub>O 0.001g, MnSO<sub>4</sub>.H<sub>2</sub>O 0.001g, Agar 15g and sterile distilled water 1ltr. The colonies were stored dry at 20°C. If they were to be required for use again they could be revived by streaking on Starch-

Casein agar and incubated at room temperature for 7 days (Bhagabati *et al.*, 2008).

**Antimicrobial activity evaluation:** Antimicrobial activity of the secondary metabolites was determined by use of Kirby Bauer disk diffusion method. One week old broth cultures of the actinomycetes were used with sterile paper disks soaked in the cultures for 30 minutes were used to inoculate actinomycetes on media seeded with pathogenic bacteria under test at 37°C for 24 hours.

## RESULTS

Halo zones were observed after 24hrs of incubation while those with resistance, there was significant growth of test organisms (Table 2).

**Table2.:** Results for sensitivity test.

Isolates.	Kp	Sf	Ec	Ss	Sa	Bs	St
Is.1		-	-	-	-	+	+
Is.2	-	-	-	-	-	+	+
Is.3	-	-	-	-	-	+	+
Is.4	-	-	-	-	-	+	-
Is.5	-	+	-	-	-	+	+
Is.6	+	+	-	-	-	+	+
Is.7	-	+	-	-	-	-	+
Is.8	-	-	-	-	+	+	+
Is.9	-	-	+	+	+	+	+
Is.10	-	+	+	+	+	+	+
Is.11	+	-	-	-	+	+	+
Is.12	-	+	-	-	+	-	+
Is.13	-	-	+	+	+	+	+
Is.14	-	+	-	+	+	+	+
Is.15	-	+	-	-	-	+	+

**Key:**

**Sf-** *Streptococcus feacalis*  
**Ec-** *E.coli*  
**Kp-** *Klebsiella pneumoniae*  
**Sa-** *Staphylococcus aureus*  
**Ss-** *Shigella sonei*

**St-** *Salmonella typhi*  
**Ps-** *Pseudomonas auregnosa*  
**Bs-** *Bacillus subtilis*  
 + - halo zone formed  
 - halo zone not formed

**Analysis:** Formation of halo zones (*Photograph 3a of Bacillus subtilis*) indicated bacterial growth inhibition ability. This showed that the actinomycete produced secondary metabolites that were active against the test


bacteria. Absence of halo zone was demonstrated that test bacteria were resistant to the action of the metabolite (*Photograph3b. Streptococcus feacalis*) (Jeffrey, 2008).



**Plate3a. Bacillus subtilis.**

**Plate3b. Streptococcus feacalis.**

Bar graph1 represents graphical presentation of the data.



**Key:**

- Sf- *Streptococcus feacalis*
- Ps- *Pseudomonas auregnosa*
- Ec- *E. coli*
- Bs- *Bacillus subtilis*
- Kp- *Klebsiella pnuemoniae*
- Sa- *Staphylococcus aureus*
- Ss- *Shigella sonei*

*Salmonella typhi* was susceptible to active metabolites from all the isolates apart from isolate 4. *Bacillus subtilis* all shown to be susceptible to almost all the isolates apart from isolate 7 and 12. *E.coli* was resistant to most of the isolates apart from 3 isolates 3, 9, 13 and 14. *Shigella sonnei* was also resistant to most of the isolates apart from

isolate 9, 10, 13 and 14. Isolates 9, 10, 13 and 14 proved to be a broader antibiotic spectrum as they acted against most of the isolates. Isolate 7 acted against growth of *Salmonella typhi* and *Klebsiella pneumoniae* while isolate 4 showed poor activity. It was only active for *Bacillus subtilis*.

## DISCUSSION

Evaluation for production of active secondary metabolites was done using the disk diffusion method. The 6mm disks soaked in the actinomycete broth absorbed the metabolites and carried the actinomycetes' spores as well. The actinomycetes were grown for one week thus enabling them to release their metabolites into the media. The antimicrobial metabolites are known to accumulate under stress conditions like depletion of nutrients (Hopwood, 2000). All the isolates showed positive results. This is because they were able to inhibit growth of some of the test bacteria as was indicated by the formation of a halo zone. Different metabolites from the different isolates acted against the different test bacteria while some metabolites acted against a number of test bacteria. Secondary metabolites from Is.4 were active against *B. subtilis* only while secondary metabolites from Is.1 were active against *B. subtilis* and *S. typhi*. Test bacteria showed varied responses on these metabolites by being susceptible to metabolites from one isolate and resistant to the other isolate. *Staphylococcus aureus* was susceptible to metabolites from Is.8, 9, 10, 11, 12, 13 and Is.14 while it was resistant to the other metabolites from the remaining isolates. The sizes of the halo zones varied with different metabolites. This was due to differences in the diffusion rates of these metabolites through the agar. These rates of diffusion were dependent on the diffusion and solubility properties of the metabolite. Faster diffusing metabolites formed larger halo zones compared to slow diffusing metabolites (Cheesbrough, 2006). Size of molecular weight of the compounds may also have had effect on the diffusion rates. Compounds with low molecular weights diffused more compared to metabolites with a large molecular weight thus formed larger halo zones (Dougherty and Pucci, 2012). The metabolites are known to diffuse in three dimensions so agar with shallow depths will have the metabolite diffusing more and thus producing large halo zones in contrast to agar with deeper depths. These factors in combination result in each combination having a unique ability in formation of

the zone of inhibition indicating susceptibility of the test bacteria to the antimicrobial substance (Jorgensen and Ferraro, 2009 and Wheat, 2001). Point at which critical mass is reached for the compounds is indicated by a sharp margin of the halo zone beyond which no inhibitory activity was evidenced. Resistance among other bacteria may be attributed to several factors among them inoculation of bacterial mass above the critical mass that overpowered the inhibition effects of the metabolites. Bedenic, Beader and Zagar in their studies to determine effect of size of inoculums on antibacterial activity of cefpirome and cefepime against *K. pneumoniae* strains known to produce SHV extended-spectrum beta-lactamases established that higher inoculums sizes rendered the strains resistant to the two test antibiotics (Bedenic, Beader and Zagar, 2001). There is also the possibility that these bacteria might have acquired the resistant genes from the neighboring resistant bacterial cells in the previous environment (Jorgensen and Ferraro, 2009 and Canton, 2008). The reason for different sensitivity between Gram positive and Gram negative bacteria could be ascribed to the morphological differences between these microorganisms, Gram negative bacteria having an outer polysaccharide membrane carrying the structural lipopolysaccharide components. This makes the cell wall impermeable to lipophilic solutes (Pandey *et al.*, 2011). The Gram positive bacteria would be more susceptible due to having only an outer peptidoglycan layer which is not an effective permeability barrier (Davies and Davies, 2010). Susceptibility might also have been contributed by several factors. Synthesis of cell walls, protein, and membrane of the test bacteria might have been inhibited. There might have been blockage of nucleotide synthesis and membrane function. This resulted to no division and death of bacterial cells. Interpretation of susceptibility or resistance is based on the presence or absence of zones of inhibition surrounding the disks (Jorgensen and Ferraro, 2009).

## CONCLUSION

Research findings from this study show promising results especially for the multi-drug resistant strains. The challenge of drug resistance has become a public health concern and by these findings, it dictates a constant need

for more discovering of novel antimicrobial drugs and explorations. This solely depends on the results of research findings on antibiotics with much emphasis on actinomycetes.

## ACKNOWLEDGEMENT

The author wishes to thank all members of staff of Department of Plant and Microbial Sciences Kenyatta University for their support and technical input. Special

thanks Dr. Kibera and Dr. Maingi for their academic advice.

## REFERENCES

- Barrios-Gonzalez, J., Fernandez, J.F & Tomasini, A. (2003). Microbial Secondary Metabolites Production and Strain Improvement. *Indian Journal of Biotechnology* 2:322-333
- Bedenic, B., Beader, N & Zagar, Z. (2001). Effect of Inoculum Size on the Antibacterial Activity of Cefpirome and Cefepime against *Klebsiella pneumoniae* Strains Producing SHV Extended-spectrum Beta-lactamases. *Journal of Clinical Microbiology and Infection* 7(11):626-635.
- Canton, R. (2009). Antibiotic Resistance Genes from the Environment: A Perspective through Newly identified Antibiotic Resistance Mechanisms in the Clinical Setting. *European Society of Clinical Microbiology and Infectious Diseases* 15(1):20–25.
- Ceylan, O., Okmen, G & Ugur, A. (2008). Isolation of Soil *Streptomyces* as Sources Antibiotics Active against Antibiotic-resistant Bacteria. *EurAsian Journal of BioSciences* 2:73-82.
- Cheesbrough, M. (2006). *Direct Laboratory Practice in Tropical Countries*, Vol.2. New York: Cambridge University Press.
- Cummins, C.S. and Harris, H. (1956). The Chemical Composition of the Cell wall in some Gram-positive Bacteria and its Possible Value as a Taxonomic Character. *Journal of General microbiology* 14(3):583-600.
- Davies, J & Davies, D. (2010). Origins and Evolution of Antibiotic Resistance. *Molecular and Molecular Biology Reviews* 74(3):417-433.
- Dougherty, J.T & Pucci, M.J. (2012). *Antibiotic Discovery and Development*. New York: Springer.
- Harvey, R.A. (2007). *Microbiology*. Baltimore: Lippincott Williams & Wilkins.
- Hopwood, D.A., Buttner, M.J., Kieser, T & Charter, K.F. (2000). Antibiotic Production by *Streptomyces*. *Practical Streptomyces Genetics* 1:1-42.
- Jeffery, L.S.H. (2011). Prescreening of Bioactivities from Actinomycetes Isolated from Forest Peat Soil of Sarawak. *J. Trop. Agric. And Fd.* 39(2):245-253.
- Jeffrey, L.S.H. (2008). Isolation, Characterization and Identification of Actinomycetes from Agricultural Soils at Semongok, Sarawak. *African Journal of Biotechnology* 7(20):3697-3702.
- Jorgensen, J.H & Ferraro, M.J. (2009). Antimicrobial Susceptibility Testing: A Review of General Principles and Contemporary Practices. *Clinical Infectious Diseases* 49(11):1749-1755.
- Kekuda T.R.P., Shobha, K.S & Onkarappa, R. (2010). Fascinating Diversity and Potent Biological Activities of Actinomycete Metabolites. *Journal of Pharmacy Research* 3(2):250-256.
- Kumar, S.K. (2001). *Actinomycetes of an Indian Mangrove (Pichavaram) Environment: An Inventory*. Ph.D. thesis, Annamalai University, India, 91 pp.
- Oskay, M., Same, A., Azeri, C. (2004). Antibacterial Activity of some Actinomycetes Isolated from Farming Soils of Turkey. *African Journal of Biotechnology* 3(9): 441-446.
- Pandey, A., Ali, I., butola, K.S., Chatterji, T & Singh, V. (2011). Isolation and Characterization of Actinomycetes From Soil Evaluation of Antibacterial Activities of Actinomycetes Against Pathogens. *International Journal of Applied Biology and Pharmaceutical Technology* 2(4):384-392
- Sacramenti, D.R., Coelho, R.R.R., Wigg, D.M., Linhares, L.T.F.L., Dos Santos, M.G.M., Semedo, L.T.A.S & Silva, A.J.R. (2004). Antimicrobial and Antiviral Activities of an Actinomycete (*Streptomyces* sp.) Isolated from a Brazilian Tropical Forest Soil. *World Journal of Microbiology and Biotechnology* 20: 225-229.
- Sumpradit, N., Chongtrakul, P., Anuwong, K., Pumtong, S., Kongsomboon., Butdeemee., Khonlormyati,

- J., Chomyong, S., Tongyoung, P., Losirawat, S., Seesuk, P., Suwanwaree, P & Tangcharoensathien, V. (2012). Antibiotics Smart Use: A Workable Model for Promoting the Rational Use of Medicines in Thailand. *Bull World Health Organization* 90:905-913.
- Wheat, P.F. (2001). History and Development of Antimicrobial Susceptibility Testing Methodology. *Journal of Antimicrobial Chemotherapy* 48(1):1-4.