

Soil mite densities from central Ivory Coast

Julien Kouadio N'Dri¹ and Henri Marc André^{2,1}

¹ Biodiversity Research Centre, Earth and Life Institute, Université Catholique de Louvain, Place Croix du Sud 4, B-1348 Louvain-la-Neuve, Belgium

² Musée royal de l'Afrique centrale, Leuvensesteenweg 13, B-3080 Tervuren, Belgium

Corresponding author email: ndri_jk@yahoo.fr

Keywords: soil mites, forest, savannah, densities.


1 SUMMARY

Four sites, the Lamto savannah, the Oumé primary forest and Oumé teak plantation (Sudanese domain) and the Taï primary forest (Guinean domain) were sampled twice (in the rainy and dry season) in Ivory Coast. During this study three hypothesis were investigated: (1) soil mite densities vary with habitat type and season; (2) soil mite densities are affected by soil physico-chemical parameters; and (3) soil mite densities vary with depth (vertical distribution) and along transects (horizontal distribution). After a 1-week extraction in Berlese-Tullgren funnels, mite densities were higher during the rainy season than during the dry season. Despite the site and the season, density generally decreased from the litter to the deep layers despite the appearance of a bimodal distribution in some sites. The seasonal effect was more marked in top soils. In spite of the season, the same density succession was observed: Oumé forest - Lamto savannah - Oumé teaks - Taï forest. Major taxa Oribatida and Gamasida decreased with the depth in all sites and in all seasons. Contrary to what is observed in temperate areas, the Soil Depth₅₀ indicated that the study of top soils may be sufficient to describe the soil mite densities in the Tropics. Physico-chemical parameters such as water content and apparent density influenced the vertical mite distribution.

2 INTRODUCTION

Soils are complex open systems in which biosphere, lithosphere, atmosphere and hydrosphere come together (Ibáñez and Boixadera, 2002). The implications of this complex three-dimensional milieu are of fundamental importance for a very diverse biota (Coleman *et al.*, 2004). Soil biological diversity is sometimes linked to its functioning (Lavelle, 1996; Andrén and Balandreau, 1999; Fitter *et al.*, 2005; Bardgett, 2008). Generally, the soil fauna, especially mites, is considered to play a key role in the soil functioning. It contributes to the decomposition process and nutrient mineralization, mixture and fragmentation of soil particles and regulation of micro-organisms, (Seastedt, 1984; Lavelle and Spain, 1991; Coleman *et al.*, 2004; Gobat *et al.*, 2004; Bardgett,

2005; Illig *et al.*, 2010). Despite several calls (Ghabbour, 1984; André *et al.*, 1992), the literature on soil microarthropods from tropical and equatorial Africa remains meager. In Africa, soil microarthropods abundance is highly variable depending on the vegetation, the habitat fragmentation and mesological factors (Badejo, 1990, 1994; Noti *et al.*, 1996; Badejo and Tian, 1999; Badejo and Ola-Adams, 2000; Noti *et al.*, 2003); although all these studies are based on topsoil and disregard mineral soils. A strong exception is offered by the publications of Athias *et al.* (1975) and Athias (1975, 1976) who studied mites from Lamto down to 40 cm. Her approach is however limited to a single site, the Lamto savannah in Ivory Coast colonized mostly by a herbaceous stratum.



This second study in Ivory Coast is original as soil mite densities were estimated in topsoil and mineral soils in different sites selected in distinct ecological zones. Most sites benefited from a special protection status (primary forest, savannah). In contrast, a teak plantation was chosen near a protected primary forest. The

following hypotheses were specifically explored: (1) soil mite densities varies with habitat type and season; (2) soil mite densities are affected by soil physico-chemical parameters; (3) soil mite densities varies with depth (vertical distribution) and along transects (horizontal distribution).

3 MATERIAL AND METHODS

3.1 Study areas: Four sites, all located in arboreal areas of Ivory Coast, were sampled in 2008. The first site (Lamto) is in a transition zone between forest and savannah in the Sudanese domain, the second and third sites (Oumé) are also located in the Sudanese domain and the last one (Taï) pertains to the Guinean domain. In general, Ivory Coast is characterized by a tropical climate of the Sudanese and Guinean type, with four alternating seasons (High dry season: December to March; great rainy season: April to July; short dry season in August and short rainy season: September to November).

3.1.1 Lamto savannah (LAS): The first site is the 2500-ha Lamto reserve (Coordinates: 6°13' N, 5°02' W; altitude: 125 m ASL). The Ecology Station is located 153 km north-west of Abidjan; its history with ecological studies is provided by Vuattoux *et al.*, (2006). The climate (intertropical humid), reviewed by Le Roux (2006), is characterized by a mean annual temperature of 27°C and a mean yearly rainfall of 1211 mm (data from Lamto Geophysical Station). The dry season occurs in December and January. Broadly defined, the savannah is perceived as a continuous layer of graminoids (grasses and sedges) and a discontinuous layer of trees and shrubs (Solbrig *et al.*, 1996). In Lamto, all facies were dominated by tall palm trees (*Borassus aethiopum*), pretty regularly distributed (Barot, 1999). Sampling was conducted in an unburned area already prospected by Athias *et al.*,

(1975) and Athias (1975, 1976) and now dominated by *Chromolaena odorata* (Asteraceae) (Fig. 1A).

Soils were ferralsols (FAO classification) with a very low organic matter and nitrogen content (1% and 0.5% respectively) (Delmas, 1967; Riou, 1974; Mordelet *et al.*, 1996).

3.1.2 Oumé primary forest (OPF): The second site is located in Oumé, mid-west Ivory Coast, precisely in the village of Goulikao (6°31' N, 5°30' W) at 200 m asl (Fig. 1B). The climate regime is subequatorial (Attic climate) (Monnier, 1983). Average temperature and relative humidity are 26°C and 85%, respectively (Avenard *et al.*, 1971; Bongoua, 2002). The rainfall pattern is of bimodal type, with two dry seasons and one rainy season. Yearly average rainfall is around 1 275 mm, with peaks reaching 2 015 mm and 1 386 mm, in June and in September, respectively. Oumé is characterized by a mesophile-type vegetation, or a semi-deciduous forest (Monnier, 1983). Vegetation is very dense and even luxuriant (Fig. 1B). Man-made activities are very weak and limited to some tracks. The undergrowth is also dense with lianas and dead wood. *Griffonia simplicifolia* (Caesalpiniaceae), *Marantochloa leucantha* (Marantaceae), *Anthiaris toxicaria* (Moraceae) are some trees found in OPF. Ferralitic soils are regularly distributed and cover all areas (Assié *et al.*, 2008).





Figure 1: The four sampling sites. LAS with *Chromolaena odorata* in foreground (A); discontinuous woody stratum in OPF (B); no herbaceous stratum and no undergrowth in OTK (C); weak light penetration in TPF (D); (Photo N'Dri, 2008, in a dry season).

3.1.3 Oumé teak plantation (OTK): The third site is also located in Oumé, near the previous forest. It is composed of even-aged teaks (*Tectona grandis*) planted in 1994 and represented the site influenced by the disturbance (Man-made activities such as clearing, cutting and associated goings-on are very high). Most conspicuous is the absence of undergrowth and herbaceous stratum; the ground is bare (Fig. 1C).

3.1.4 Taï primary forest (TPF): The Taï National Park is the largest remaining forest (450 000 ha) in West Africa and the last largest island of the original Upper Guinean forest that once reached from Ghana to Guinea-Bissau. Sampling was carried out at about 200 m North of the central research station (approximate coordinates: 5°45' N 7°07' W, altitude 150-200 m asl). The climate is of subequatorial type, humid all the year round (Kouadio, 2006; OIPR, 2006). The yearly average rainfall over the period

1993-2002 was 1 853 mm, with a mean annual temperature of 25.35°C. The Taï National Park is dominated by *Eremospatha macrocarpa* and *Diospyros mannii* forests. The flora was estimated to 1 231 species (Adou *et al.*, 2005). The Taï forest, although not layered, is definitely structured and the light penetration to the undergrowth is weak (Alexandre, 1982a, b) (Fig. 1D). Highly desaturated ferrallitic soils and hydromorphic soils cover almost all areas (Avenard *et al.*, 1971; Moreau, 1983).

3.2 Sampling method: Mite data were obtained from 1 080 soil cores, *i.e.* 8 series of 15 sets scattered over a 14-m transect, homogeneous from a floristic point of view. Each set determined a “biological” profile representing the mite vertical distribution and was composed of 9 cores, 1 from litter (organic layer; variable height) and 8 from mineral soil taken at regular depths below the litter (0-5 cm ; 5-10 cm ; 10-



15 cm ; 15-20 cm ; 20-25 cm ; 25-30 cm ; 30-35 cm ; 35-40 cm). Series of 15 sets (the litter sample and eight 48-cm³ samples) were repeated in the 4 sites and 2 seasons (dry season in January to March 2008; rainy season from September to October 2008 for most sites, December 2008 for TPF). In other words, the sampling design was as follows: 4 sites x 15 sub samples collected along each transect x 9 layers x 2 campaigns = 1080 units. Because of the difficulties in doing controlled field experiments at regional scales, this sampling protocol takes greater advantage of natural field observations and avoids pseudo replicates (Hargrov and Pickering, 1992; Kozlov and Hurlbert, 2006; Knezevic, 2009). A steel corer (Ø 3.5 cm) was used to take samples vertically. It was not possible to take samples more deeply as the corer met rocky materials.

3.3 Microarthropod extraction: Microarthropods were extracted using the Berlese–Tullgren funnels. Light bulbs (25 W), situated 15 cm above the intact samples, and were lit immediately after the beginning of the extraction process. Extraction lasted 1 week (after 7 days, no arthropod was captured in preliminary samples). The mean top temperature in the extraction cabinet did not exceed 39°C. This method was chosen after preliminary studies.

3.4 Mite identification: Microarthropods were sorted in Petri dishes using a dissecting microscope. Mites were mounted in lactic acid or in Hoyer medium for identification under a light microscope with phase contrast. Major taxa (Oribatida, Gamasida, Actinedida and Acaridida) are defined in the traditional sense used in soil zoology and available in Krantz (1978) and Dindal (1990).

3.5 Physico-chemical parameters: In organic soils, the sample depth corresponds to the mean litter height (variable depending on the site and season); in mineral soils, it corresponds to the mean layer depth plus the mean litter height. Other physico-chemical parameters were estimated from seven cores adjacent to the sample itself, five cores were mixed together to measure the pH and chemical analyses as recommended by Anderson and Ingram (1993), one core was reserved for the water content and the last one for the apparent or bulk density. Soil water content and apparent density of the 960 mineral soil samples were estimated after 48 h drying at 105°C (Baize, 1988; Duchaufour, 1991).

The pH (water) (Baize, 1988; Duchaufour, 1991) was estimated with a pH meter (HANNA) after calibration at the “Laboratoire d’Écologie du sol”. Other chemical analyses for extreme depths (0-5 cm and 35-40 cm) were conducted by the “Centre Provincial de l’Agriculture et de la Ruralité” located in La Hulpe (Belgium). To measure organic and total carbon, the carbon was oxidised to CO₂ by heating the soil to at least 900°C in a flow of oxygen-containing gas according to ISO Standard 10 694. Total Nitrogen was determined by dry combustion, according to ISO Standard 13 878. Soil organic matter (SOM) was estimated through the formula $C_{org} \times 1.7$ as made by Noti *et al.*, (2003).

3.6 Data analysis: Mite abundances, *i.e.* the number of individuals/sample, were transformed into density. As soils are three-dimensional bodies of the earth's crust with properties than can vary over different spatial and temporal scales in every direction (Ibáñez and Boixadera, 2002; Castrignanò *et al.*, 2004; Blum, 2006), densities were expressed in number of mites/dm³ (as in Table 1). Nevertheless, soils are generally investigated only in horizontal dimensions and densities and comparisons with previous studies were often made in mites/m² (as in Table 4). The SE (Standard Error) associated with mite densities was high and distribution did not follow the normality. Therefore, non parametric (Mann-Whitney) tests were used to compare mite densities in the different layers, as one moves deeper into the soil. Multiple tests may be difficult to interpret “because if we go on testing long enough we will inevitably find something which is *significant*” (Bland and Altman, 1989). However, the sequential Bonferroni was not utilized as encouraged by Moran (2003) and García (2004). Comparisons between sites and seasons were stopped whenever tests were not significant. As explained by André *et al.*, (2002), we estimated the soil depth in which 50 or 90% of arthropods were living (respectively the Soil Depth₅₀ and Soil Depth₉₀, abbreviated as SD₅₀ and SD₉₀).

Soil physico-chemical means were compared using a one-way ANOVA test. Rank Pearson correlation was preferred to study the relationship between mite density and soil physico-chemical parameters. All tests were conducted using STATISTICA 7.1 software package (Statsoft, Tulsa, USA).




Table 1: Mite density (mean individuals/dm³ ± SE) in litter (N = 15), mineral soils (0-40 cm, N = 120), and all samples (N = 135, global density) in the four sites and two seasons.

	LAS		OPF		OTK		TPF					
Dry season												
Litter	52	±	50	56	±	42	32	±	38	50	±	65
0-40 cm	13	±	27	38	±	51	8	±	27	5	±	16
Global density	18	±	32	42	±	51	12	±	30	11	±	30
Rainy season												
Litter	117	±	154	233	±	116	92	±	73	83	±	96
0-40 cm	38	±	117	51	±	103	14	±	28	11	±	26
Global density	48	±	123	75	±	119	21	±	43	16	±	46


Table 2: Seasonal effect on mite density (individuals/dm³) in different layers (N = 15) and the entire profile (N = 9) of the four sites (P-values of Mann-Whitney tests).

	LAS	OPF	OTK	TPF
Litter	0.310	0.000 ***	0.010 **	0.141
0-5 cm	0.044 *	0.003 **	0.245	0.351
5-10 cm	0.237	0.431	0.078	0.507
Profile	0.427	0.566	0.122	0.200

Significant at levels 0.05 (*), 0.01 (**) and 0.001 (***).

Table 3: Effects of sites on mite density (individuals/dm³) in different layers (N = 15) and the entire profile (N = 9) in different seasons (P-values of Mann-Whitney tests).

	DRY SEASON			RAINY SEASON		
	LAS	OPF	OTK	LAS	OPF	OTK
Litter						
OPF	0.632			0.004 **		
OTK	0.272	0.110		0.709	0.001 ***	
TPF	0.917	0.494	0.468	0.983	0.000 ***	0.576
0-5 cm						
OPF	0.272			0.089		
OTK	0.373	0.082		0.044 *	0.000 ***	
TPF	0.093	0.012 *	0.619	0.026 *	0.000 ***	0.272
5-10 cm						
OPF	0.019 *			0.548		
OTK	0.229	0.001 ***		0.407	0.034 *	
TPF	0.351	0.002 **	0.772	0.178	0.018 *	0.468
10-15 cm						
OPF	0.017 *			0.300		
OTK	0.221	0.001 ***		0.340	0.020 *	
TPF	0.120	0.000 ***	0.756	0.967	0.245	0.245
15-20 cm						
OPF	0.065			0.034 *		
OTK	0.507	0.002 **		0.756	0.054	




TPF	0.507	0.002	**	1.000	0.756	0.054	1.000
20-25 cm							
OPF	0.002	**			0.135		
OTK	1.000	0.002	**		0.590	0.075	
TPF	0.787	0.002	**	0.787	0.885	0.147	0.772
25-30 cm							
OPF	0.013	*					
OTK	0.983	0.017	*				
TPF	0.756	0.028	*	0.787			
25-30 cm							
OPF	0.384						
OTK	0.330	0.054					
TPF	0.330	0.054		1.000			
Profile							
OPF	0.013	*			0.251		
OTK	0.158	0.007	**		0.596	0.085	
TPF	0.133	0.007	**	0.596	0.402	0.027	* 0.566

Significant at levels 0.05 (*), 0.01 (**) and 0.001 (***).

Table 4: Density (mean individuals/m²) in litter and mineral soils (0-40 cm) of the major groups of mites in the four sites and two seasons. Totals and percentages are also given.


	Litter	0-5	5-10	10-15	15-20	20-25	25-30	30-35	35-40	Total	%
Dry season											
LAS											
Gamasida	1 109	555	69	555	139	69	0	416	69	2 981	35
Actinedida	208	139	0	139	69	0	0	0	0	555	6
Oribatida	2 080	1 733	485	139	139	69	208	139	0	4 992	59
Acaridida	0	0	0	0	0	0	0	0	0	0	0
Total	3 397	2 426	555	832	347	139	208	555	69	8 527	100
%	40	28	7	10	4	2	2	6	1	100	
OPF											
Gamasida	1 941	1 941	832	971	277	485	693	485	416	8 042	39
Actinedida	139	69	0	208	0	0	0	0	69	485	2
Oribatida	3 120	2 288	971	1941	832	1456	485	347	485	11 924	58
Acaridida	0	0	0	0	0	0	0	0	0	0	0
Total	5 200	4 298	1803	3120	1109	1941	1179	832	971	20 451	100
%	25	21	10	15	5	9	6	4	5	100	
OTK											
Gamasida	1 387	1 387	139	69	0	69	69	0	139	3 258	57
Actinedida	139	0	0	0	0	0	0	0	0	139	2
Oribatida	832	555	0	69	69	0	0	69	0	1 595	28
Acaridida	139	277	0	0	0	69	277	0	0	763	13
Total	2 496	2 218	139	139	69	139	347	69	139	5 754	100
%	44	39	2	2	1	2	7	1	2	100	
TPF											
Gamasida	277	208	69	0	0	0	277	69	0	901	17



Actinedida	69	0	0	0	0	0	0	0	0	69	1
Oribatida	2 842	1 040	139	0	69	139	69	0	0	4 298	82
Acaridida	0	0	0	0	0	0	0	0	0	0	0
Total	3 189	1 248	208	0	69	139	347	69	0	5 269	100
%	61	24	4	0	1	2	7	1	0	100	
Rainy season											
LAS											
Gamasida	5 338	5 962	1 941	416	69	208	208	139	69	14 351	65
Actinedida	139	139	69	0	0	0	0	0	69	416	2
Oribatida	1 317	2 218	555	139	69	69	0	69	69	4 506	20
Acaridida	69	69	0	0	0	0	0	2 634	0	2 773	13
Total	6 863	8 389	2 565	555	139	277	208	2 842	208	22 046	100
%	31	38	12	3	1	1	1	12	1	100	
OPF											
Gamasida	6 794	6 239	1 525	485	832	277	69	69	277	16 569	48
Actinedida	0	0	69	0	0	0	69	0	0	139	0
Oribatida	6 378	4 576	1 040	277	1 525	485	901	485	69	15 737	46
Acaridida	624	832	277	0	69	69	69	0	0	1 941	6
Total	13 796	11 647	2 912	763	2 426	832	1 109	555	347	34 386	100
%	40	34	8	2	7	2	4	2	1	100	
OTK											
Gamasida	1 525	1 040	347	139	69	69	277	485	69	4 021	44
Actinedida	69	0	0	0	0	208	208	555	0	1 040	11
Oribatida	1 941	1 387	277	69	139	0	139	69	0	4 021	44
Acaridida	0	0	0	0	0	0	0	0	0	0	0
Total	3 536	2 426	624	208	208	277	624	1 109	69	9 082	100
%	39	27	7	2	2	3	7	12	1	100	
TPF											
Gamasida	693	485	347	69	69	277	69	139	0	2 149	30
Actinedida	277	69	0	0	0	0	69	0	0	416	6
Oribatida	1 941	1 525	208	277	139	69	208	0	0	4 368	61
Acaridida	0	69	0	139	0	0	0	0	0	208	3
Total	2 912	2 149	555	485	208	347	347	139	0	7 141	100
%	41	30	8	7	3	5	5	1	0	100	

Table 5: Soil Depth₅₀ and Soil Depth₉₀ (in centimeters) estimated for the four sites and in the two seasons.

	LAS	OPF	OTK	TPF
Dry season				
SD ₅₀	5.1	13.4	4.7	2.2
SD ₉₀	26.6	35.6	29.7	25.0
Rainy season				
SD ₅₀	5.6	4.4	3.5	3.0
SD ₉₀	34.5	22.2	32.4	22.6



4 RESULTS

4.1 Mites density: In all sites and seasons, the mite density generally decreased from the litter to the deep layers (Fig. 2) and was higher in organic than in mineral soils (Table 1). The highest densities were observed in OPF, especially in the rainy season (233 ± 116 mites/dm³ in the litter, 233 ± 183 mites/dm³ in the 0-5 cm layer). No mites were captured in the deepest layers of TPF, and also in the 10-15 cm layer in dry season of TPF. Generally, the lowest densities were found in OTK and TPF, and in the dry season rather than in the rainy season (except in TPF where the climate is humid all the year round). The ratio of immature mite (1.65%) was weak and did not affect significantly the densities.

Whatever the season, the same global density succession OPF-LAS-OTK-TPF was observed (Table 1). The SE values associated with mean densities were high as cited above and not illustrated in Fig. 2. Tests comparing profile densities between seasons were not significant (Table 2). However, seasons significantly affected the mite densities observed in litter (OPF and OTK) and in the 0-5 cm layer (LAS and OPF). Deeper, the season effect was not significant. In the dry season, the mean mite density observed in litter was not significantly different between the four sites (Table 3). In mineral soils and in the entire profile however, there was a significant difference in densities surveyed in OPF compared to the other sites.

In the rainy season, the opposite trend was observed in litter: OPF was significantly different from other sites. The same trend was found in mineral soils, but less deep than in the dry season. In addition, densities recorded in the 0-5 cm layer in LAS were significantly greater (8389 ind/m²) than those in OTK (2426 ind/m²) and TPF (2149 ind/m²) (Table 4). Profile densities differed significantly only in OPF and TPF.

4.2 Major taxa: Oribatida dominated in most sites (LAS: 59%, OPF: 58%, TPF: 82%) in the dry season and in TPF (61%) only in the rainy season. In other cases, Gamasida represented the dominant group (LAS: 65%, OPF: 48%, OTK: 44% in the dry season and OTK: 57% in the rainy season) (Table 4). Whatever the season Oribatida and Gamasida were the majors dominant group, respectively in TPF and OTK. Actinedida were a minor group, with a maximum of 11% in OTK in the rainy season. Acaridida constituted a second minor group, with a maximum of 13% in OTK in the dry season. The major group, Oribatida, Gamasida and Acaridida were significantly ($N = 36$; $p \geq 0.05$) different between the

four sites in the dry season. However in the rainy season, major group are not different in the four site ($N = 36$; $p \geq 0.05$).


4.3 SD₅₀ and SD₉₀: All SD₅₀ values, except OPF in the dry season, were less than the litter height (Table 5). SD₉₀ was around 30 cm.

4.4 Physico-chemical parameters: In spite of the site and the season, apparent density increased significantly ($N = 120$, $P < 0.05$) from the 0-5 to 35-40 cm layers (Fig. 3A, B). Apart from the 10-15 cm layer in the dry season, the apparent density also differed significantly ($N = 60$, $P < 0.05$) between sites. Mean values observed in each site were close to each other whatever the season (LAS: 1.04 in the dry season vs. 0.90 in the rainy season; OPF: 1.22 vs. 1.04; OTK: 1.03 vs. 1.05; TPF: 1.06 vs. 1.12).

Contrary to apparent density, water content (Fig. 3C, D) significantly decreased with layer depth in the two seasons ($N = 120$, $P < 0.05$), except in LAS in the dry season ($P = 0.928$). Water contents were also significantly different when sites were compared ($N = 60$, $P < 0.05$). Mean values were: LAS: 8.08% in the dry season vs. 14.63% in the rainy season; OPF: 7.03 vs. 15.29%; OTK: 12.51 vs. 18.21%; TPF: 19.48 vs. 14.81%). Note that the values obtained in TPF were in the reverse order than in other sites.

TPF soils were acidic and the pH values differed significantly from those observed in other sites ($N = 120$, $P < 0.05$; Fig. 3E, F). Beside, the pH gradient in TPF was reverse from that noticed elsewhere: the deeper the layer, the greater the pH. The pH values obtained in the layers of the three remaining sites differed significantly, except in OTK in the rainy season ($P = 0.054$). Mean values were similar in the two seasons: LAS: 6.52 in the dry season vs. 6.48 in the rainy season; OPF: 7.20 vs. 7.49; OTK: 7.32 vs. 7.34; TPF: 6.01 vs. 5.88). Whatever the season, C_{org} contents were identical with C_{tot} contents, except in the 0-5 cm layer in OTK during the rainy season. Carbon and nitrogen values were greater in 0-5 cm layer than in deeper soils (Table 6). These values were significantly different ($P < 0.05$) between sites except for the 35-40 cm layer in the dry season.

4.5 Mite density and physico-chemical parameters: In OPF during the dry season only, litter height and pH were significantly correlated to the mite density (Table 7). Except in OTK in the rainy season, there was a significant and negative correlation between depth and mite density. Apart from TPF in the dry season and OTK in the rainy




season, apparent density and water content were also significantly correlated to the mite density. Last, there was a significant correlation between mite density and

C_{tot} , N_{tot} , C/N and SOM in the 0-5 cm layer observed in LAS in the rainy season.

Table 6: Mean and SE values of chemical parameters measured in the two extreme layers of mineral soils in the four sites and two seasons.

	LAS			OPF			OTK			TPF		
DRY SEASON												
0-5 cm												
C _{org} (g/kg)	9.60	±	0.68	26.60	±	2.69	16.80	±	1.39	23.60	±	2.77
C _{tot} (%)	0.96	±	0.07	2.66	±	0.27	1.68	±	0.14	2.36	±	0.28
N _{tot} (%)	0.09	±	0.01	0.27	±	0.03	0.18	±	0.02	0.19	±	0.02
C/N	10.64	±	0.26	9.69	±	0.23	9.35	±	0.05	12.49	±	0.51
SOM (g/kg)	16.32	±	1.15	45.22	±	4.58	28.56	±	2.37	40.12	±	4.71
35-40 cm												
C _{org} (g/kg)	5.30	±	0.00	6.18	±	0.72	5.44	±	0.14	7.20	±	0.97
C _{tot} (%)	0.53	±	0.00	0.62	±	0.07	0.54	±	0.01	0.72	±	0.10
N _{tot} (%)	0.03	±	0.00	0.06	±	0.01	0.05	±	0.01	0.07	±	0.01
C/N	15.90	±	1.08	10.93	±	1.00	12.31	±	0.94	10.85	±	0.68
SOM (g/kg)	9.01	±	0.00	10.51	±	1.22	9.25	±	0.24	12.24	±	1.65
RAINY SEASON												
0-5 cm												
C _{org} (g/kg)	11.80	±	1.59	36.80	±	6.30	16.60	±	0.87	17.80	±	1.39
C _{tot} (%)	1.18	±	0.16	3.68	±	0.63	1.72	±	0.14	1.78	±	0.14
N _{tot} (%)	0.11	±	0.02	0.37	±	0.06	0.18	±	0.01	0.14	±	0.01
C/N	10.99	±	0.11	9.92	±	0.33	9.50	±	0.28	12.49	±	0.40
SOM (g/kg)	20.06	±	2.71	62.56	±	10.70	28.22	±	1.48	30.26	±	2.37
35-40 cm												
C _{org} (g/kg)	5.30	±	0.00	5.30	±	0.00	5.44	±	0.14	6.00	±	0.00
C _{tot} (%)	0.53	±	0.00	0.53	±	0.00	0.54	±	0.01	0.60	±	0.00
N _{tot} (%)	0.03	±	0.00	0.04	±	0.00	0.04	±	0.01	0.05	±	0.00
C/N	17.67	±	0.00	13.60	±	1.14	12.95	±	1.35	12.00	±	0.00
SOM (g/kg)	9.01	±	0.00	9.01	±	0.00	9.25	±	0.24	10.20	±	0.00

**Table 7:** Pearson correlation between mite density (mean individuals/dm³) and mesological variables in the four sites and two seasons. App. density = apparent density.

	LAS				OPF				OTK				TPF			
	<i>N</i>	<i>r</i>	<i>P</i>		<i>N</i>	<i>r</i>	<i>P</i>		<i>N</i>	<i>r</i>	<i>P</i>		<i>N</i>	<i>r</i>	<i>P</i>	
DRY SEASON																
Entire profile																
Litter height	15	-0.09	0.751		15	0.53	0.038	*	15	0.18	0.518		15	-0.33	0.227	
App. density	8	-0.86	0.005	**	8	-0.88	0.003	**	8	-0.70	0.048	*	8	-0.70	0.051	
Water content	8	0.86	0.005	**	8	0.78	0.020	*	8	0.77	0.023	*	8	0.70	0.052	
pH	8	0.70	0.051		8	0.81	0.014	*	8	0.62	0.099		8	-0.18	0.667	
Depth	9	-0.81	0.007	**	9	-0.77	0.015	*	9	-0.68	0.041	*	9	-0.71	0.029	*
0-5 cm																
C _{tot}	5	0.62	0.262		5	-0.25	0.679		5	-0.39	0.510		5	0.74	0.149	
N _{tot}	5	0.63	0.254		5	-0.12	0.842		5	-0.39	0.510		5	0.67	0.208	
C / N	5	0.24	0.694		5	-0.67	0.215		5	0.31	0.603		5	0.12	0.846	
SOM	5	0.62	0.262		5	-0.25	0.670		5	-0.39	0.510		5	0.74	0.149	
35-40 cm																
C _{tot}	5	-	-		5	-0.06	0.920		5	-	-		5	-	-	
N _{tot}	5	-	-		5	0.21	0.723		5	-	-		5	-	-	
C / N	5	-	-		5	-0.58	0.296		5	-	-		5	-	-	
SOM	5	-	-		5	-0.06	0.920		5	-	-		5	-	-	
RAINY SEASON																
Entire profile																
Litter height	15	-0.21	0.435		15	-0.29	0.283		15	-0.13	0.629		15	0.33	0.283	
App. density	8	-0.74	0.032	*	8	-0.89	0.002	**	8	-0.52	0.186		8	-0.78	0.020	*
Water content	8	0.89	0.002	**	8	0.94	0.001	***	8	0.63	0.091		8	0.78	0.021	*
pH	8	0.69	0.054		8	0.4	0.316		8	0.46	0.248		8	-0.59	0.115	
Depth	9	-0.68	0.043	*	9	-0.8	0.008	**	9	-0.66	0.050		9	-0.76	0.015	*
0-5 cm																
C _{tot}	5	0.93	0.020	*	5	0.39	0.512		5	0.49	0.397		5	0.24	0.687	
N _{tot}	5	0.93	0.020	*	5	0.25	0.681		5	0.37	0.533		5	0.30	0.614	
C / N	5	-0.88	0.045	*	5	0.79	0.107		5	0.67	0.213		5	-0.41	0.491	
SOM	5	0.93	0.020	*	5	0.39	0.512		5	0.38	0.522		5	0.24	0.687	
35-40 cm																
C _{tot}	5	-	-		5	-	-		5	-	-		5	-	-	
N _{tot}	5	-	-		5	0.00	1.000		5	-	-		5	-	-	
C / N	5	-	-		5	-0.07	0.901		5	-	-		5	-	-	
SOM	5	-	-		5	-	-		5	-	-		5	-	-	

Significant at levels 0.05 (*), 0.01 (**) and 0.001 (***). -: spurious correlation





Figure 2: Mite density (individuals/dm³) in the litter (L followed by the average height in cm) and mineral soil (depths of layers) in the four sites and during the two seasons. LAS: (A, B); OPF: (C, D); OTK: (E, F); TPF: (G, H); dry season: (A, C, E, G); rainy season: (B, D, F, H).






Figure 3: Physico-chemical parameters (mean values and SE) in the four sites and during the two seasons. Apparent density: (A, B); water content: (C, D); pH: (E, F); dry season: (A, C, E); rainy season: (B, D, F).



5 DISCUSSION

5.1 Efficiency of the Berlese-Tullgren funnels: The efficiency rate of traditional extraction methods is especially weak and the most frequently used extraction methods have a mean efficiency of 40%, a rough overestimation advanced by André *et al.*, (2002). The absolute efficiency of the Berlese-Tullgren funnels would vary, depending on the authors, from 26% (Forsslund, 1948) to 7% (André *et al.*, 2002) and the Berlese-Tullgren funnels are also selective with respect to their efficiency for certain taxa (variable taxonomic and functional efficiency).

The Berlese-Tullgren funnels were supposed little efficient in Ivory Coast as the ratio of mite immatures is especially weak (less than 2%) and the rate of Actinedida varied from 2 to 6%. Last, the comparison of tolerance zones, critical and lethal temperatures and temperature preferences made by Wallwork (1960) between Oribatida from Ghana (West Africa) and Michigan (North America) showed major differences which plea for a weak efficiency of funnels. Nevertheless, the Berlese-Tullgren funnels were retained for their low cost and feasibility in Ivory Coast. All densities published in this paper rely on abundances in berlesates and are thus rough estimations.


5.2 Comparison with Lamto results: Mite densities observed in LAS soils, 9 000 to 22 000 mites/m² compared to a yearly 28 000 mites/m² (Athias, 1975) are similar as well as the extraction method, the Berlese-Tullgren funnels, and the sample size. Water contents, C_{tot}, N_{tot} and SOM also remain the same compared with Athias's data, even if the herbaceous vegetation has changed over time. Contrary to Athias' (1976) conclusions, water content seems to govern the mite density observed in Lamto. Another discrepancy with previous results concerns the importance of major taxa: Gamasida from 35 to 65% of mites vs. 16% in Athias (1975) and Actinedida from 2 to 6% vs. 39% in Athias (1975). This difference within major taxa could be attributed to changes observed in the vegetation structure. Indeed, the plot sampled by Athias was dominated by herbaceous stratum (Poaceae). In contrast our plot was characterized by the marked presence of *Chromolaena odorata*. The litter of this plant decomposed very quickly in organic matter. This idea was confirmed by the positive correlation between mite density and the chemical parameters (C_{tot}, N_{tot}, SOM).

5.3 Comparison with other data from Africa:

Whatever the site, mite density is greater in the rainy than in the dry season. A similar trend is reported by Badejo and Tian (1999) in southwest Nigeria. Such a trend could be attributed to a low soil moisture regime associated to a high soil temperature (Badejo, 1990; Badejo *et al.*, 1998; Badejo and Tian, 1999). During the rainy season, the litter accumulated throughout the dry season starts disappearing gradually due to decomposition. Maldague (1961) and Noti *et al.*, (1996) showed that mites were particularly abundant in African forests. The highest densities of soil mites observed in OPF might be explained by the heterogeneity of vegetation and ensuing microhabitats. Such a hypothesis is supported by the positive correlation with the litter height estimated in the dry season. However, the oribatid density observed in OPF is higher in the rainy than in the dry season, which contradicts the results published by Noti *et al.*, (1996)

At the same sampling depth (7.5 cm) plus litter and whatever the season, mite density in OPF (dry season: 10 399 individus/m²; rainy season: 16 899 individus/m²) is greater than that reported in Nigeria (1 792 individus/m²; 8 500-10 000 individus/m²) respectively by Badejo and Tian (1999) and Adeduntan (2009). Mite densities in OPF remain however very weak compared to results by Noti *et al.*, (1996) (rainy season: 30 700 individus/m²; dry season: 37 200 individus/m²) and limited to adult oribatids. In Congo Democratic Republic, they are even less than values recorded in the Namib Desert by André *et al.*, (1997), although these authors were used carbon tetrachloride flotation method.

Contrary to expectation and despite the absence of disturbance lowest densities are reported from TPF where the climate is humid all the year round. Highly desaturated ferrallitic soils and hydromorphic soils cover almost all areas (Avenard *et al.*, 1971; Moreau 1983). Soil type could be an obstacle to the mites' development in this zone. These results agree with recent observations on spiders (R. Jocqué, pers. obs.) and Myriapoda (D. Van den Spiegel, pers. obs.) on the same site (soil acid), but require further investigations. Low densities are also reported in OTK, where there is no undergrowth and the litter is monospecific. Oribatid densities in OTK however are clearly higher than reported by Badejo and Ola-Adams (2000) in a similar habitat. The same major group dominant, mesostigmatid mite was also observed by Badejo and



Ola-Adams (2000), in spite of the soil sampled was acid (pH: 5.18) and low in organic matter (3.43 %). The low density of mites in the teak plantation could be explained by the disturbance and soil compaction during the establishment of plants. The soil bulk density was higher in the upper layer and in this case, limited the water infiltration.

A seasonal effect is visible only in litter (OTK and OPF) and in the upper layer of mineral soil (LAS and OPF). These results suggest the presence of surface species sensitive to a seasonal effect, which agrees with Badejo's (1990) findings.

5.4 Influences of physico-chemical parameters:

Whatever the season and whatever the site, the litter height and the pH little affect the mite vertical distribution. However, parameters as soil apparent density (or porosity alike), and water content play a key role in determining the mite density (Noti, 1991;

Ducarme *et al.*, 2004). In contrast, C_{tot} , N_{tot} , C/N and SOM were correlated with mite density only in LAS and seemed to have a minor role.

5.5 Soil Depth₅₀: Most values of SD₅₀ are below 5.6 cm and clearly indicate that a study of topsoil may suffice to describe the soil mite densities from Ivory Coast. The only exception is OPF in the dry season (SD₅₀ = 13.4 cm). This study conclusion is conflicting to that reported by Athias (1975) who claimed that, contrary to what is observed in temperate areas; microarthropods from tropics have a deep distribution. This study results confirm the findings by Adis *et al.*, (1987) and Illig *et al.*, (2010) that most arthropods live in the top 3.5 cm in Manaus (Brazil) but are opposite to André's *et al.* (2002) estimation of 11 cm for the SD₅₀ in temperate forests. However this vertical distribution could be affected by an altitude variation between the study sites (Illig *et al.*, 2010).

6 CONCLUSION

The different results reveal that the density of mites is higher when the site is undisturbed, as is the case with Oumé primary forest. However this factor alone could not explain the weak density observed in others sites. Other parameters such as soil properties, the speed of litter decomposition, and ecosystem productivity could greatly influence the mite

abundance. The relatively high density of mites in Lamto savannah will be the response of site protection and its transition to woodland. This study results also suggest a variable distribution of soil mite density. This has to be supported by biological diversity studies as proposed by André *et al.* (1992).


7 ACKNOWLEDGMENTS

Special thanks to T. Hance. Thanks also to E. J. Tondoh, R. Jocqué, Y. Samyn, and D. Van den Spiegel for advices and grants. Financial support from


CSM-BGBD project, FRS/UCL, GTI/IRScNB, ABIC/MRAC is gratefully acknowledged.

8 REFERENCES


- Adeduntan SA: 2009. Diversity and abundance of soil mesofauna and microbial population in South-Western Nigeria. *African Journal of Plant Science* 3(9): 210-216.
- Adis J, De Moraes JW. and Ribeiro EF: 1987. Vertical distribution and abundance of arthropods in the soil of a neotropical secondary forest during the dry season. *Trop. Ecol.* 28: 174-181.
- Adou YA, Blom EC, Dengueadhe KTS, Van Rompaey RSAR, N'Guessan EK, Wittebolle G. and Bongers F: 2005. Diversité floristique et végétation dans le Parc National de Taï, Côte-d'Ivoire. Tropenbos International, Wageningen, The Netherlands.
- Alexandre DY: 1982a. Pénétration de la lumière au niveau du sous-bois d'une forêt dense tropicale. *Ann. Sci. forest* 39: 419-438.
- Alexandre DY: 1982b. Strata in tropical rain-forest at Taï (Ivory Coast). *Tropical Rain Forest, The Leeds Symposium*, Leeds Philosophical and Literary Society: 15-24.
- Anderson JM. and Ingram JSI (Eds.): 1993. *Tropical Soil Biology and Fertility. A Handbook of methods* (2nd ed.). CAB International, Wallingford, UK.
- André HM, Ducarme X. and Lebrun Ph: 2002. Soil biodiversity: myth, reality or conning? *Oikos* 96: 3-24.
- André HM, Lebrun Ph. and Noti M-I: 1992. Editorial. *Biodiversity in Africa: a plea for more data.* *J. Afr. Zool.* 106: 3-15.



- André HM, Noti M-I. and Jacobson KM: 1997. The soil microarthropods of the Namib Desert: A patchy mosaic. *J. Afr. Zool.* 111: 499-517.
- Andrén O. and Balandreau J: 1999. Biodiversity and soil functioning—from black box to can of worms? *Appl. Soil Ecol.* 13: 105–108.
- Assié KH, Angui P. and Tamia AJ: 2008. Effets de la mise en culture et des contraintes naturelles sur quelques propriétés physiques d'un sol ferrallitique au Centre Ouest de la Côte d'Ivoire: conséquences sur la dégradation des sols. *European Journal of Scientific Research* 23: 149-166.
- Athias F: 1975. Données complémentaires sur l'abondance et la distribution des microarthropodes de la savane de Lamto (Côte d'Ivoire). *Bull. Mus. Natl Hist. Nat.* 308: 1-28.
- Athias F: 1976. Recherche sur les microarthropodes du sol de la savane de Lamto (Côte-d'Ivoire). *Ann. Univ. Abidjan, sér. E, Ecologie* 9: 193-271.
- Athias F, Josens G. and Lavelle P: 1975. Traits généraux du peuplement animal endogé de la savane de Lamto (Côte-d'Ivoire). In Vanek J. (Ed.). *Progress in Soil Zoology*, pp. 375-388. Academia Publishing House, Prague, Czech Republic.
- Avenard J-M: 1971. Aspect de la géomorphologie. In *Le milieu naturel de la Côte-d'Ivoire, Mémoires ORSTOM* 50: 70-72.
- Badejo MA: 1990. Seasonal abundance of soil mites (Acarina) in two contrasting environments. *Biotropica* 22: 382-390.
- Badejo MA: 1994. Effect of accidental fire on soil mite density in a forest reserve in Nigeria. *Experimental and Applied Acarology* 18: 703-710.
- Badejo MA, Nathaniel TI. and Tian G: 1998. Abundance of springtails (Collembola) under four agroforestry tree species with contrasting litter quality. *Biol. Fertil. Soils* 27: 15-20.
- Badejo MA. and Ola-Adams BA: 2000. Abundance and diversity of soil mites of fragmented habitats in a biosphere reserve in southern Nigeria. *Pesq. agropec. Brasília* 11: 2121-2128.
- Badejo MA. and Tian G: 1999. Abundance of soil mites under four agroforestry tree species with contrasting litter quality. *Biol. Fertil. Soils* 30: 107-112.
- Baize D: 1988. Guide des analyses courantes en pédologie. INRA Éditions, Paris, France.
- Balogh J. and Balogh P: 1992. The Oribatid Mites Genera of the World (vol. 1 and 2). the Hungarian National Museum Press, Budapest, Hungary.
- Bardgett RD: 2005. *The Biology of Soil. A Community and Ecosystem Approach.* Oxford University Press, Oxford, UK.
- Bardgett RD: 2008. The consequences of soil biodiversity for ecosystem function. Presented to Soils 2008, Conference Massey University, Palmerston North, New Zealand, 1-5 December 2008 (Unpublished).
- Barot S: 1999. Interactions entre répartition spatiale, hétérogénéité environnementale et démographie : cas du palmier Rônier dans une savane humide de Côte d'Ivoire. *Ecologie*. PhD Dissertation. Université de Paris 6, France.
- Blum WEH: 2006. The future of soil science. In Hartemink AE. (Ed.). *The Future of Soil Science*, pp. 16-18. IUSS International Union of Soil Sciences, Wageningen, The Netherlands.
- Bongoua AJ: 2002. Caractérisation de l'état de fertilité des jachères plantées en légumineuses fixatrices d'azote dans la région d'Oumé. DEA Dissertation, Université de Cocody-Abidjan, Côte d'Ivoire.
- Castrignanò A, De Giorgio D, Fornaro F. and Vonella AV: 2004. 3D spatial variation of soil impedance as affected by soil tillage. In Raine SR, Biggs AJW, Menzies NW, Freebairn DM, Tolmie PE (Eds). *ISCO 2004 - 13th International Soil Conservation Organisation Conference - Brisbane, 4-9th July 2004. Conserving Soil and Water for Society: Sharing Solutions. Paper 744*, pp. 1-5, Australian Society of Soil Science Inc., Brisbane, Australia.
- Coleman DC, Crossley Jr DA. and Hendrix PF: 2004. *Fundamentals of Soil Ecology* (2nd ed.). Academic Press, Burlington, U.S.A.
- Delmas J: 1967. Recherches d'écosystèmes dans les savanes de Lamto (Côte d'Ivoire). Premier aperçu sur les sols et leur valeur agronomique. *La terre et la vie* 21: 216-227.
- Dindal DL (Ed.): 1990. *Soil Biology Guide*. Wiley J. and Sons, New York, U.S.A.



- Ducarme X, André HM, Wauthy G. and Lebrun Ph: 2004. Comparison of endogeic and cave communities: microarthropod diversity and mite species richness. *European Journal of Soil Biology* 40: 129-138.
- Duchaufour P: 1991. *Abrégés de pédologie. Sol, végétation, environnement* (3^{ème} éd.). Masson, Paris, France.
- Fitter AH, Gilligan CA, Hollingworth K, Kleczkowski A, Twyman RM, Pitchford JW. and the members of the nerc soil biodiversity programme: 2005. Biodiversity and ecosystem function in soil. *Functional Ecology* 19: 369–377.
- Forsslund K-H: 1948. Något om insamlingsmetodiken vid markfaunaundersökningar. *Medd. Statens Skogforskningsinst* 37(7): 1–22.
- García LV: 2004. Escaping the Bonferroni iron claw in ecological studies. *Oikos* 105: 657-663.
- Ghabbour SI: 1984. Literature on soil fauna of Africa needed! *Biology International*, 9: 20.
- Gobat J-M, Aragno M. and Matthey W: 2004. *The Living Soil. Fundamentals of Soil Science and Soil Biology*. Science Publishers, Enfield (NH), USA, Plymouth, UK.
- Hargrove WW. and Pickering J: 1992. Pseudoreplication: *a sine qua non* for regional ecology. *Landscape Ecology* 6: 251-258.
- Ibáñez JJ. and Boixadera J: 2002. The search for a new paradigm in pedology: a driving force for new approaches to soil classification. In Micheli F, Nachtergaele RJ, Jones A, Montanarella L. (Eds). *Soil Classification 2001*. European Soil Bureau - Research Report 7: 93-110.
- Illig J, Norton AR, Scheu S. and Maraun M: 2010. Density and community structure of soil land bark-dwelling microarthropods along an altitudinal gradient in a tropical montane rainforest. *Exp Appl Acarol*. doi:10.1007/s10493-010-9348-x.
- Kouadio KKF: 2006. Analyse du système de biomonitoring du Parc National de Taï. École Supérieure d'Agronomie (ESA), Institut National Polytechnique Houphouët Boigny de Yamoussoukro (Côte d'Ivoire) (MSc Dissertation).
- Kozlov MV, Hurlbert SH: 2006. Pseudoreplication, chatter, and the international nature of science: A response to DV Tatarnikov. *J. Fund. Biol. (Moscow)* 67: 145-152.
- Knezevic A: 2009. Diagnosing and avoiding pseudoreplication. *StatNews* 75: 1-2.
- Krantz GW: 1978. *A Manual of Acarology* (2nd ed.). Oregon State University Bookstores, Corvallis, U.S.A.
- Lavelle P: 1996. Diversity of soil fauna and ecosystem function. *Biology International* 33: 3-16.
- Lavelle P. and Spain A: 1991. *Soil Ecology*. Kluwer, Dordrecht, The Netherlands
- Le Roux X: 2006. Climate. In Lamto. Structure, Functioning, and Dynamics of a Savanna Ecosystem. Abbadie L, Gignoux J, Le Roux X, Lepage M. (Eds). Lamto. Structure, Functioning, and Dynamics of a Savanna Ecosystem, pp. 25-44. Springer Verlag, New York, USA.
- Maldague M: 1961. Relation entre le couvert végétal et la microfaune, leur importance dans la conservation biologiques des sols tropicaux. *Publ. Inst. Natn. Étude agron. Congo, Sér. scientifique* 90: 1-122.
- Monnier Y: 1983. Végétation. In Vennetier P. (Ed.), *Atlas de la Côte d'Ivoire* (2nd ed.), p. 17, Jeune Afrique, Paris, France.
- Moran MD: 2003. Arguments for rejecting the sequential Bonferroni in ecological studies. *Oikos* 100: 403-405.
- Mordelet P, Barot S. and Abbadie L. 1996. Root foraging strategies and soil patchiness in a humid savanna. *Plant Soil* 182: 171–176.
- Moreau R: 1983. Sur l'origine d'éléments d'aspects charbonneux observés dans les sols de la région de Taï. Office de la Recherche Scientifique et Technique d'Outre-Mer, Centre d'Adiopodoumé (Côte d'Ivoire), 6 pp. (unpublished).
- NotM-I: 1991. Etude biocénétique des peuplements des acariens du sol du Haut-Shaba dans la région de Lubumbashi, Zaïre. PhD Dissertation. Université Catholique de Louvain.
- Noti M-I, André HM. and Dufrêne M: 1996. Soil oribatid mite communities (Acari: Oribatida) from high Shaba (Zaïre) in relation to vegetation. *Applied Soil Ecology* 5: 81-96.
- Noti M-I, André HM, Ducarme X. and Lebrun Ph: 2003. Diversity of soil oribatid mites (Acari: Oribatida) from high Katanga (Democratic



- Republic of Congo): a multiscale and multifactor approach. *Biodiversity and Conservation* 12: 767-785.
- OIPR: 2006. Plan d'aménagement et de gestion du Parc National de Taï. Office Ivoirien des Parcs et Réserves, Ministère de l'Environnement, des Eaux et Forêts, République de Côte d'Ivoire
- Riou G: 1974. Les sols de la savane de Lamto. *In* Analyse d'un écosystème tropical humide : la savane de Lamto (Côte d'Ivoire). Les facteurs physiques du milieu, Bull. de liaison des chercheurs de Lamto 3: 3-45.
- Seastedt TR: 1984. The role of microarthropods in decomposition and mineralization processes. *Annu. Rev. Entomol* 29: 25-46.
- Solbrig OT, Medina E. and Silva JF: 1996. Biodiversity and tropical savanna properties : A global view. *In* Mooney HA, Cushman JH, Medina E, Sala OE, Schulze ED. (Eds.). *Functional Roles of Biodiversity: A Global Perspective*, pp. 185-211, John Wiley and Sons Ltd, New York, U.S.A.
- Vuattoux R, Konaté S, Abbadie L, Barot S, Gignoux J. and Lahoreau G. 2006. History of the Lamto Ecology Station and Ecological Studies at Lamto. *In* Abbadie L, Gignoux J, Le Roux X, Lepage M. (Eds). *Lamto. Structure, Functioning, and Dynamics of a Savanna Ecosystem*, pp. 1-12. Springer Verlag, New York, USA.
- Wallwork JA: 1960. Observations on the behaviour of some oribatid mites in experimentally-controlled temperature gradients. *Proc. Zool. Soc. London* 135: 619-29.