



# First record of the avian malaria vector *Cs. longiareolata* (Diptera: Culicidae) for the Southeast of Algeria



<https://doi.org/10.35759/JABs.154.2>

## ABSTRACT

**Objectives:** The control strategies of mosquitos like *Culiseta longiareolata* were developed gradually. A study of the bio-ecology of *Culiseta longiareolata* was considered as the first step to fight against it.

**Methodology and results:** Larvae and adult collections were carried out from different habitats using standard dipping methods from the region of Biskra (Algerian Southeast) (34° 51' 00" north, 5° 44' 00" east). For five years during 2009-2013 in four sites and the investigation was bi-monthly for any site. Overall, 792 specimens of mosquitoes were identified among the 44677 specimens that were collected. *Cs. longiareolata* presented in the second order with 27.59 % in numerical terms after *Culex pipiens* 28.65 %, *Aedes caspius* 12.97 %, *Anopheles multicolor* 7.22 %, *Anopheles sergentii* 2.84 %. Other 17 species were presented with low rates. This species was collected from all types of habitats (dominant or periodic stagnant water, with or no vegetation, clear or turbid, sunny ,or noisy sites, filled with fresh or brackish water, clean or polluted), throughout the year with different abundances. The Phenology of *Cs. longiareolata* proves that it is distributed in ten months annually, compared with other species. The sexual dimorphism was clear across the morphometric characters of both sexes. The densities were variables for this species when marked two picks, for the month of May (> 40 individual/stroke for larvae, > 32 individual/stroke for an adult), for the month of November (> 27 individual/stroke for larvae, > 20 individual/stroke for an adult).

**Conclusion and application of findings:** More, *Cs. longiareolata* cohabited with all species of different genera *Aedes*, *Culex*, and *Anopheles*.

**Keywords:** Culicidae, bioecology, *Culiseta longiareolata*, Biskra, breeding sites.

## INTRODUCTION

Avian malaria and various arboviruses are very interesting diseases on the veterinary and ecological side; of which *Cs. longiareolata* (Macquart, 1838) is the main vector (Seidel et al., 2013). It is also the vector involved in the

transmission of brucellosis, avian influenza, West Nile encephalitis, and Malta fever (Maslov, 1967). This species is multivoltine with continuous development in hot countries with wide distribution, is present in the south of the Palearctic region; it is

distributed in the Mediterranean region, Europe, and Asia (Van Pletzen et al., 1989; Azari et al., 2011, Minar 1991). Actually, there is a contradiction with many authors mentioning that this species attacks human and animal, but the birds are only the principal host in North Africa (Al-Jaran & Katbeh-Bader, 2001; Maslov et al., 1989; Roubaud & Colas-Belcour, 1989). The breeding sites of this species are of very different types; they may be permanent or temporary, shaded or sunny, filled with fresh or brackish water, clean or polluted (Schaffner, 2001). *Cs. longiareolata*, is a widely distributed species in Algeria (Lounaci, 2003; Berchi, 2000; Hamaidia, 2004), and particularly in the southern part of the country (Clastrier & Senevet, 1961; Mengri et al., 1984; Merabti, 2016; Merabti et al., 2017). To fight

against insects which have vectorial capacities is a major concern for scientists. A good understanding of the ecology of species is classified as a first step to control these vectors. In a period of five years from January 2009 to December 2013 and in four different sites of the region of Biskra (Southeast of Algeria), the study was done on the ecological factors impact on the spatio-temporal distribution of *Cs. longiareolata*. The biometric study of twenty-six morphological characters of the two sexes of this species can show his systematic aspect. In the Biskra (Algeria) region, like everywhere in the state, the direct effect of the water in the natural environment creates a biotope in organic matter, in nitrogen and phosphate products, favourable to the proliferation of mosquitoes.

**MATERIAL AND METHODS**

**Presentation of the study area:** A preliminary survey was carried out in natural and urban areas in the region of Biskra (34°51'N 5°44'E), in the South-East of Algeria, which allowed us to inventory about ten larval sites

where Culicidifauna was harvested. Four different collection sites were chosen in this study area. Samples were taken twice a week in six cottages for each site. The description of the study sites is in table 01:

**Table 1:** Natural characteristics of the study sites.

Site Characteristics	Geographic coordinates	Breeding sites type	Sampling Technique
01: CHETMA	34°50'53.02"N and 5°48'01.42" E	Swamp, puddle, irrigation basin, runoff	Butterfly net, Dipping methods
02: BISKRA	34°50'14.66"N and 5°44'37.93" E	Valley, Basin water	Butterfly net, Dipping methods, hand catch
03: SIDI OKBA	34°45'22.16"N and 5°54'04.64" E	Dam water	Butterfly net, Dipping methods
04: BORDJ BEN AZOUZ	34°41'43.33"N and 5°21'52.79" E	Irrigation ditches, freshwater or brackish pond	Butterfly net, Dipping methods

**Sampling method:** Mosquito development stages were sampled at 24 breeding habitats twice a week using Dipping method (Papierok et al., 1975), and butterfly net during all the month in the five years of study. Sampled specimens were preserved in vials containing ethanol (75%). In addition, mosquitoes were sampled manually with handheld aspirators and tubes. Each vessel was

carefully marked with information about the breeding site. The specimens sampled (pre-imaginal stages and imagines) were identified morphologically using the keys of Brunhes et al., 2000). Data Analysis: The graphical representations used in this study were prepared using Exel (2007), Image Tools v.4.2 and Statistix v8.0.

**RESULTS**

**The numeric importance of *Cs. longiareolata*:** In the five years of prospection, in the number of collected specimens two species *Culex pipiens* (the domestic mosquitoes) that have a large local spread and of *Cs.*

*longiareolata* were marked as important. The number reported is varied between 467 and 79 individuals for the first and second season, 2009 and 2013 consecutively. The last season had a decreased number of the

collected specimen maybe due to climatic factors like temperature and precipitation or because of the very hot year of 2012 in the region of Biskra, when the temperature has an important value about 50 to 58 C°

(NMO, 2014). The third species was *Aedes ochleratatus caspius*, but it has a decreased number compared to *Cx.pipiens* and *Cs.longiareolata*. The other species with a decreased number did not arrive at 200 individuals.




Figure 1: The numeric importance of *Cs. longiareolata* with other species collected

**Breeding sites :** All positive breeding sites were cohabited by all developmental stages of *Cs. longiareolata*. During the five years of study, this species has a large capacity to occupy the different breeding sites, with nature different (permanent or temporary), and of different types (Lagoon depression, grassy marsh, drainage ditch, traditional basin water, irrigation basin, marsh and rejects sites). Were harvested in almost all types of breeding sites with different localized plant species (Green Algae, *Phragmites comunis*, *Phoenicia dactylifora*, *Cynodon dactylon*).

**Temporal distribution and phenology of *Cs.longiareolata*:** The table 2 proved the monthly


repartition of the sampling species, the *Cs.longiareolata* was observed in ten months in the year, in the five years of sampling, compared to other species, when it was differentiated remarked in the breeding sites from two to four months like *Cs.ochroptera*, *Cs. subochrea*, *Culex laticinctus*, *Culex theileri*, *Anopheles sergentii*, and *Aedes vexans*. Some species have a large monthly repartition when it observed from five to seven months like *Anopheles multicolor* and *Culex modestus*. This repartition prouve the large ecological valence of the *Cs.longiareolata* compared to others species.

Table 2: Monthly repartition of *Cs.longiareolata* with other species

Species	Months	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
---------	--------	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

<i>Ae. caspius</i>																				
<i>Ae. annulipes</i>																				
<i>Ae. vexans</i>																				
<i>Ae. dorsalis</i>																				
<i>An. multicolor</i>																				
<i>An. sergentii</i>																				
<i>Cx. modestus</i>																				
<i>Cx. theileri</i>																				
<i>Cx. laticinctus</i>																				
<i>Cx. torrentium</i>																				
<i>Cs. Longiareolata</i>																				
<i>Cs. Annulata</i>																				
<i>Cs. Subochrea</i>																				
<i>Cs. Ochroptera</i>																				

Ae: Aedes; An: Anopheles; Cx: Culex; Cs: Culiseta.



L; liter, 1 to 12: months.

**Figure 2:** Monthly density of larvae (2009-2013)


**Monthly density according to Dipping method:**

According to the method of Dipping, which consists of sampling larvae and pupae without repetition in several places at each breeding site using a 1-liter ladle. We can remark that the density of the *Cs. longiareolata* larvae were differently from month to another. We can remark that the density is very important for the month of May when the density has touched the value of 40 Larvae/ladle, then the three months of March, April, and November, when density was from 25 to 32 Larvae/ladle. The density was observed with a low average compared to previous months (5 to 20 larvae/ladle) (Figure 2).

**Density of adults according to the Butterfly net:**

For the adult's density was determined by Butterfly net used during our sampling. The values were very important for the three months of April, May, and November, when it accessed 20 adults/net. This observation was revealed maybe to the climatic conditions: temperature and the wind for example (Figure 3).

According to Figure 2 and 3, we observed that the density of larvae of *Cs. longiareolata* was higher than the density marked for adults while the 12 months in the five years of prospection.




L: liter, 1 to 12: onths

**Figure 3:** Monthly density of adults (2009-2013)


**Morphometric study:** The measurements of descriptive criteria selected of both sexes were illustrated in Figures.4 and 5. It's presented on histogram compared the mean measurement and standard deviation of 26 criteria related to females and males respectively. Those

entire measurements proved the sexual dimorphism between both sexes when the females are smaller than the males. These morphometric characteristics present a specificity of our species in the south of our country.



WL: Wing length; WW: Wing Width; Ant:Antenna; Plp: Palp; Pbs: proboscis; T: Tarsus.

**Figure 4:** Histogram presentation of mean measurements of *Cs. longiareolata* females



WL: Wing length; WW: Wing Width; Ant:Antenna; Plp: Palp; Pbs: proboscis; T: Tarsus

**Figure 5:** Histogram presentation of the mean measurement of *Cs. longiareolata* males.

Figure 5, presented on the histogram compared the mean measurement and standard deviation of 26 criteria related to males. The analysis of variance (ANOVA) between both sexes, with the twenty-six characters measurements, proves that there is a high significant difference ( $F^{182,4}=39.3$ ;  $p=0.000$ ). According to the study of Senevet, 1936; Clastrier & Senevet 1961; Merabti (2016); Merabti et al., 2016; Bebbi, 2004, The *Cs. longiareolata* was sampled from many sites from the Sahara regions of Algeria (Biskra, Oues Souf, Laghouat, Ghardia, Ouargla, and the south of the region of Djelfa). According to Brunhes et al., (1999) this species is multivoltine, stenogamous and autogenic, its role as a vector of human parasitosis can only be reduced. According to Systematic Catalogue of Culicidae, 2011; Becker et al., 2010, this species is very spread in many countries in Europe and Africa. Tine-Djebbar et al., 2011 announced that the *Cs. longiareolata* and *Cx. pipiens* are the two most remarkable mosquito species in Algeria. The female also bites man and pets, but has not been implicated in the transmission of human parasitosis (Senevet & Andarelli, 1959; Brunhes et al., 1999). According to Boulkenafet, (2006), the eggs of *Cs. longiareolata* grouped in the basket are cylindro-conical, carries about 50 to 400 eggs. For the year 2012, the temperatures recorded in our region of the study experienced very high values, which exceeded 50 ° c. At

this point, we have reported the almost total absence of *Cs. longiareolata* populations in the breeding sites. This remark can interfere with the limits of its ecological valence, which is limited by high-temperature values. This can interpret the oviposition behaviour of the female (Spencer et al., 2002; Stav et al., 2000). In recent years, our species has been found in urban areas, so the stinging behaviour has become anthropogenic rather than just zoophilic. The impacts of spatially heterogeneous environmental and ecological factors on mosquito population dynamics are complex and poorly understood for many species like, Particularly for zoonotic mosquito-borne diseases like *Cs. longiareolata*. This specie was sampled from different breeding sites (artificial or artificial), with stagnant or running water, rich or without aquatic flora, exposed or covered by the sun, pure and polluted water. This species were sampled as well the field and urban sites, but surely observed it did not represent a nuisance to man as it is in the animal's blood like Dove and Rabbit (Medlock et al., 2005; Kampen et al., 2013; Merabti et al., 2017) Brunhes et al., 2001 was confirmed that this species is a widely distributed species. It is found in artificial and natural deposits (Rioux, 1958). During our surveys, we found it in all the cottages and during most of the year (except the month of July) in the permanent lodgings with fresh water poor or rich in vegetation. Lounaci (2003) reported

its existence in the Reghaia marsh deposit, in the deposits of the El Harrach Agricultural Institute and in the El-Alia barn. Agoun (1996) and Berchi (2000) reported its presence in different types of deposits in Constantine. Hamaidia (2004) reports its existence in polluted deposits, permanent settlements with stagnant water rich or poor in vegetation and in temporary dwellings with stagnant or current water with or without vegetation, in the regions of Tébessa and Souk-Ahras. Zittra et al, (2014) confirmed in his research that the developmental stages of *Cs. longiareolata* were exclusively detected in one artificial water storage container and not at natural breeding sites. These findings confirm existing data, which indicate that *Cs. longiareolata* mainly favors man-made and predator-free artificial water containers and temporary waters with low predation risk for their offspring (Stav et al. 1999, Kiflawi et al. 2003). This study results were according to Zittra et al (2014) that the larvae and pupae of *Cs. longiareolata* shared their habitat with specimens of the *Culex pipiens* complex and *Cs. Annulata*, but we noticed that this species can cohabit its breeding sites with other species like *Cs.subochrea*, *Cs.ochroptera*, *Cx.modestus*, *Cx.theileri*, *Cx.laticinctus*, *Cx. torrentium*, *An.sergentii*, *An.multicolor*, *Ae.vexans*,

#### CONCLUSION AND APPLICATION OF RESULTS

The present study was done during five years of prospection of the Saharan region of Algeria and precisely in the southeastern part of the country. Monitoring the population of *Cs.longiareolata* allowed us to release some information. This species is very widespread throughout the year and in all types of breeding sites. It can withstand very low temperatures, and high temperature, which does not exceed 50 ° C. In

#### ACKNOWLEDGMENT

This study is the fruit of five years prospection; we are especially thankful to all collaborators (biology department, laboratory, and personals to contribute to

#### REFERENCES

Agoun M, 1996. Contribution à l'étude d'un inventaire systématique des moustiques (Culicidae-Diptera) de la région de Constantine et ses abords. Mém. DES, Université de Constantine : 26p.

Al-Jaran, T. K., & Katbeh-Bader, A. M, 2001. Laboratory studies on the biology of *Culiseta longiareolata* (Macquart)(Diptera: Culicidae). *Aquatic Insects*, 23(1): 11-22.

*Ae.caspius*. We determined on the results obtained, the monthly distribution of *Cs.longiareolata* during the five years of study. This distribution has shown that this species is omnipresent all the year round, and which does not support either low or high temperature. With regard to quantitative results, the densities of this larval or adult species undergo spatio-temporal variations comparable to those reported by several authors in the different regions. It appears that spatiotemporal variations in densities cannot be attributed solely to fluctuations in the mesological parameters (environmental parameters), but is also due to the nature of the habitat that is close to the sampling station. In fact, this last factor would influence firstly the spatio-temporal distribution of Culicid species on the one hand, and the physicochemical factors of water, which can play the role of a limiting factor (Brunhes et al., 1999; Senevet & Andarelli, 1960, Rioux, 1958). The study of morphological characteristics of this species aims to distinguish the differences between the two sexes on the one hand and on the other hand to present the local ecotype of our species occupying the Saharan region of Algeria.

terms of seasonal distribution, this species occupies nine to ten months in the year and has two peaks of propagation in the two months of May and November. In these two months, the larval and imaginal densities can reach very high values. the information provided by this study can help to define ways to control this species, through its bioecology.

richness this study, A special thanks to Pr. Ralph Harbach to rework to this paper.

Azari Hamidian S, Abai MR, Arzamani K, Bakhshi H, Karami H, Ladonni H. and Harbach RE, 2011. Mosquitoes (Diptera: Culicidae) of North Khorasan Province, northeastern Iran and the zoogeographic affinities of the Iranian and Middle Asian mosquito fauna. *Journal of Entomology*, 8 (3): 217-204.

Bebba N, 2004. Etude comparative des effets d'insecticides sur les populations larvaires des Culicidae de Constantine et de Oued Righ

- (Tougourt et Djamaa).Mémoire de Magistère. Université de Mentouri. Constantine.102p.
- Berchi S, 2000. Bio écologie du *Culex pipiens* (Diptera, Culicidae) dans la région de Constantine et respectives de lutte – Thèse de Doctorat des sciences, option Entomologie. Univ. Constantine, 133 pp.
- Boulkenafet F, 2006. Contribution à l'étude de la biodiversité des Phlébotomes (Diptera : Psychodidae) et appréciation de la faune Culicidienne (Diptera : Culicidae) dans la région de Skikda. Présentation pour l'obtention du Diplôme de Magister en entomologie (option; application agronomique et médicale). 191p.
- Brunhes J, Rhaim A, Geoffroy B, Angel G and Hervy JP, 1999. Les Culicidae de l'Afrique méditerranéenne, logiciel d'identification et d'enseignement, IRD (France).
- Brunhes J, Rhaim A, Geoffroy, B, Angel G. and Hervy JP, 2000. Les moustiques de l'Afrique méditerranéenne: logiciel d'identification et d'enseignement.
- Clastrier J and Senevet G, 1961. Les moustiques du sahara central.edi Institut Pasteur d'Algérie et Laboratoire d'Entomologie der Service Antipaludique de la Délégation Générale du Gouvernement en Algerie. T. XXXZX, no 2. p25.
- D.P.A.T, 2005. Direction d'Aménagement de Territoire et de Planification. Rapport annuel sur la monographie de la wilaya de Biskra, 22p.
- Hamaidia H, 2004. Inventaire et biodiversité des Culicidae (Diptera-Nematocera) dans la région de Souk-Ahras et de Tébessa (Algérie). Thèse de Magister. Univ. Constantine.
- Kampen H, Kronefeld M, Zielke D. and Werner D, 2013. Three rarely encountered and one new Culiseta species (Diptera: Culicidae) in Germany. J Eur Mosq Control Assoc, 31, 36-39.
- Kiflawi M, Blaustein L. and Mangel M, 2003. Oviposition habitat selection by the mosquito *Cs. longiareolata* in response to the risk of predation and conspecific larval density. Ecological Entomology. 28 : 168–173.
- Lounaci Z, 2003. Biosystematics and bioecology of Culicidae (Diptera - Nematocera) in rural and agricultural environments. Thesis by Magister. INA. El Harrach.
- Maslov A VE, Ward R A and Rao PM, 1989. Blood-sucking mosquitoes of the subtribe Culisetina (Diptera, Culicidae) in world fauna (Vol. 81, No. 52003). Smithsonian Institution Libraries.
- Maslov AV, 1967. Bloodsucking Mosquitoes of the Subtribe Culisetina (Diptera, Culicidae) in World Fauna. Translation of: Krovososushchie komary podtriby Culisetina (Diptera, Culicidae) mirovo; fauny, Akademiya Nauk SSSR, Opredeliteli po Faune SSSR, Izdavaemye Zoologicheskim Institutom Akademii Nauk SSSR, Nauka Publishers, Leningrad Division, Leningrad, Amerind Publishing Co. Pvt. Ltd., New Delhi, 1989, 93: 48-100.
- Medlock JM, Snow KR. and Leach S, 2005. Potential transmission of West Nile virus in the British Isles: an ecological review of candidate mosquito bridge vectors. Medical and veterinary entomology, 19 (1): 2-21.
- Mengri M, Bessi GE and Berhaili M, 1984. Caractères Eco-Ethologiques des Culicidae et de Phlébotome ,Lutte contre ces espèces.Mimoire de fin d'étude. Ecole paramédiale de Biskra. 36 p.
- Merabti B, 2016. Identification, composition et structure des populations Culicidienne de la région de Biskra (Sud-est Algérien). Effets des facteurs écologiques sur l'abondance saisonnière. Essais de lutte (Doctoral dissertation)
- Merabti B, Lebouz, I, Adamou A E, Kouidri M. and Ouakid ML, 2017. Effects of certain natural breeding site characteristics on the distribution of Culicidae (Diptera) mosquito species in southeast Algeria. *African Entomology*, 25(2) : 506-514.
- Minar J, 1991. Family Culicidae. In: Soos A, Papp L, eds., Catalogue of Palaearctic Diptera. Volume 2. Elsevier, Amsterdam.Mosquitoes and their control. Springer, Heidelberg, Dordrecht, New York, 577 pp.
- National meteorological office, 2014. Climatological data of Biskra region (2009-2013).
- Papierok B, Croset H. and Rioux JA, 1975. Estimation de l'effectif des populations larvaires d'*Aedes cataphylla* Dyar, 1916 (Diptera, Culicidae). II. Méthode utilisant le coup de louche ou dipping. Cahiers ORSTOM, Série Entomologie Médicale et Parasitologie 13: 47–51.
- Rioux J A. 1958. Les Culicidae du « Midi » méditerranéen.Etude systématique et écologique, Ed. Paulle chevalier, Paris : 301 p.
- Roubaud PE. and Colas-Belcour J, 1933. Notes sur un culicidé méditerranéen, Theobaldia



- longiareolata Macq. Bull Soc Pathol Exot 26: 934–937.
- Schaffner F, 2001. Les Culicidae de l'Europe méditerranéenne (logiciel d'identification). Instituts des recherches pour le développement.france.2001. ISBN 2-7099-1485-9.
- Seidel B, Nowotny N, Duh D, Indra A, Hufnagl P. and Allerberger, F. 2013. First records of the thermophilic mosquito *Cs. longiareolata* (Macquart, 1838) in Austria, 2012, and in Slovenia, 2013. J Eur Mosq Control Assoc, 31: 17-20.
- Senevet G and Andarelli L, 1959. Un nouveau caractère pour la diagnose des larves de *Culex*. Arch. Inst. Pasteur, Algérie, 37 (3): 447- 461.
- Senevet G. and Andarelli L, 1960. Contribution à l'étude de la biologie des moustiques en Algérie et dans le Sahara algérien. Arch. Inst. Pasteur Algérie, t. XXXVIII (2) : 305 – 326.
- Senevet G, 1936. Notes sur les moustiques, 4-Quelques culicidés de la région de l'Aurès Algérie. *Archives de l'Institut Pasteur d'Algérie*, 14 (4) : 432-448.
- Spencer M, Blaustein L, & Cohen JE, 2002. Oviposition habitat selection by mosquitoes (*Culiseta longiareolata*) and consequences for population size. *Ecology*, 83(3) : 669-679.
- Stav G, Blaustein L. and Margalith J, 1999: Experimental evidence for predation risk sensitive oviposition by a mosquito, *Cs. longiareolata*. *Ecological Entomology*. 24,;202–207.
- Stav G, Blaustein L, & Margalit Y, 2000. Influence of nymphal *Anax imperator* (Odonata: Aeshnidae) on oviposition by the mosquito *Culiseta longiareolata* (Diptera: Culicidae) and community structure in temporary pools. *Journal of Vector Ecology*, 25 : 190-202.
- Systematic Catalog of Culicidae. 2011. Walter Reed Biosystematics Unit. ([www.mosquitocatalog.org/taxon](http://www.mosquitocatalog.org/taxon)).
- Tine-Djebbar F, Larhem AB and Soltani N, 2011. Enzyme immunoassay measurements of the molting hormone in different post-embryonic stages of two mosquito species, *Culex pipiens* and *Culiseta longiareolata*. *African Journal of Biotechnology*, 10(67): 15195-15199.
- Van Pletzen R. and Van Der Linde TDK, 1981. Studies on the biology of *Cs. longiareolata* (Macquart)(Diptera: Culicidae). *Bulletin of Entomological Research*, 71(1): 71-79.
- Zittra C, Waringer J, Werblow A, Melaun C and Fuehrer H.P, 2014. Reconfirmation of *Cs. (Allotheobaldia) longiareolata* (Macquart 1838) (Diptera: Culicidae) in Austria. The first sequence-confirmed findings in northeastern Austria. *Acta ZooBot Austria* 150/151: 17–24